

## An Asian Penis: Case Study of an Uncircumcised Organ

Penile circumcision is a routine practice for most males in the world today. The hygienic and sexual benefits have been known for many centuries. For most societies it is inconceivable to leave their males uncircumcised past puberty (the optimal age for circumcision). It is important to note that several societies still do not practice routine circumcision, most notably the Chinese. This study aims to familiarize the general public with an uncircumcised penis, its characteristics, and functioning. There are also chapters that deal with comparisons to the genitals of other circumcised Caucasian men.

### **ANATOMY**

For this study a 30-year-old, healthy Chinese male was selected. He is of average height and weight for his race (5'9", 150 lbs). His penis is, obviously, uncircumcised. Below are images and measurements of the genitals, with average Asian & Caucasian statistics for comparison:



#### Flaccid

length: 2  $\frac{3}{4}$  in (7 cm)

circumference: 3  $\frac{1}{4}$  in (8.5 cm)

#### Average Asian

3  $\frac{1}{4}$  in (8.5 cm)

3  $\frac{1}{2}$  in (9 cm)


#### Average Caucasian

4  $\frac{1}{4}$  in (11 cm)

3  $\frac{3}{4}$  in (9.5 cm)

This particular male's penis is small for his size. It is  $\frac{1}{2}$  in shorter than the average Asian and a full 1  $\frac{1}{2}$  inches shorter than the average Caucasian organ. Subject's testicles are also of smaller than average size, even by Asian standards. By comparison, the Caucasian male (far right) has much larger testes that hang about an inch lower. Also note the prominent circumcision scar behind his glans. It is rare to find adult Caucasians uncircumcised.

Asian males tend to have shorter foreskins. In adults, the glans is often slightly uncovered (as evidenced by the above subjects). The partial exposure of the glans leads to a slight decrease in smegma production. This phenomenon may be the reason why Asian males have found less reason to practice routine circumcision.



Above is the typical circumcised penis of a Caucasian male, to outline the various anatomical properties. Notice the clear circumcision scar and dry glans. This male was probably circumcised before puberty.



A detailed view of subject's flaccid penis with foreskin fully retracted. Note how the foreskin wrinkles behind the corona. Many Asians living overseas keep their foreskins retracted in this fashion during genital exposure (when urinating, in changing rooms) in an effort to hide their uncircumcised status. The fear of ridicule for having an unhygienic and unsightly organ is enough for most men to undergo a very painful adult circumcision.



Even with the foreskin retracted like this, it is easy for others to differentiate between circumcised and uncircumcised penises. The loose wrinkled foreskin behind the glans is very noticeable, with more than a passing glance. The lack of a circumcision scar (usually an inch behind the glans) is another telltale sign. If that were not enough, traces of smegma can often be found on the glans and in the folds of the foreskin. The accompanying odor can be embarrassing to the uncircumcised man when using a urinal.

The following photos illustrate the subject's penis at full erection, with average Asian and Caucasian members for comparison. Subject has a "bottle-type" shaft, where the penis narrows at the neck. Note the way subject pushes his scrotum back in order to maximize the penis length.

The circumcised Caucasian penile skin is taught (as seen below). The glans color matches the rest of the penis, unlike the moist glans of the two uncircumcised examples above it.



Erect

length: 4 ¾ in (12 cm)  
 circumference:  
 -glans: 4 ½ in (11.5 cm)  
 -shaft: 4 ¼ in (11cm)  
 -base: 4 ¾ in (12 cm)

Average Asian

5 ¼ in (13.5 cm)  
 4 ¾ in (12 cm)  
 4 ½ in (11.5 cm)  
 4 ¾ in (13 cm)

Average Caucasian

6 ¼ in (15 cm)  
 5 ¼ in (13.5 cm)  
 5 in (13 cm)  
 5 in (13 cm)

## **MASTURBATION**

It is understood that all males masturbate frequently. The sex drives in adolescent boys and adult men are extremely high. Nearly 100% of adult males masturbate with an average frequency of 6 times a week (circumcised men), compared with only 30% of adult females with an average of once a week. Given the heavy usage that the penis goes through in its lifetime, every male should take good care of its health & hygiene.

Research has found that uncircumcised males tend to masturbate more frequently than their circumcised counterparts, averaging twice daily (more than double the rate of circumcised males). There are several reasons for this. The extra nerve-endings in the foreskin contribute to the penis being more easily stimulated, thus more frequent erections. The extra sensitivity of the foreskin & moist, covered glans also promote quicker ejaculations. Premature ejaculation is almost solely the condition of uncircumcised men.

Our subject reports daily masturbation. He admits that some days he does this two or more times. This is typical for an uncircumcised male. With manual stimulation, the subject ejaculates within 2 minutes, also typical. By contrast, a circumcised Caucasian man will orgasm after 8 mins of masturbation.

The subject was observed sliding his glans in-and-out of his foreskin during masturbation. This mimics the action of sexual intercourse. This is yet another factor in the lack of stamina experienced by uncut men.

## **SEXUAL FUNCTIONING**

The time it takes for a male to climax while masturbating is a direct correlation to his stamina during intercourse with a female. The subject ejaculates within 5 mins of vaginal penetration, while the circumcised male can last up to 20 mins. This is another reason for women's preference for the circumcised organ.

Oral sex is particularly unpleasant if performed on an uncircumcised penis, unless the foreskin and glans have been thoroughly cleansed very recently. The accumulated smegma has a distinct cheesy odor and sour taste. In a recent study of 500 subjects, 92% of women polled said they would never give oral sex to an uncircumcised penis. 7% stated they would do it if the penis had just been washed. 1% would tolerate the smell. 100% preferred circumcised penises.

## **HYGIENE**

As noted previously, the primary reason for routine circumcision has been ease of hygiene. Sexual functioning is another benefit. After circumcision, the penis does not need any special attention beyond regular washing during showers. Removal of the foreskin eliminates smegma production entirely. However, in the uncircumcised penis, the foreskin secretes smegma continuously. For this reason, uncircumcised males must clean underneath the skin often.

Daily washing of the foreskin and glans is the bare minimum for the uncircumcised. Many men will need to clean twice or even three times a day, depending on the amount of smegma produced. They must also clean after each ejaculation, to prevent infections from residual semen. As uncircumcised males masturbate more often, this can be a problem for some.

Even with frequent washing, the foreskin will still retain an unmistakable odor that many women find unpleasant. What is colloquially referred to as “dick-smell”. This odor is noticeable on the underwear of almost all uncircumcised men. The only permanent solution to this problem is circumcision.

Below is an example of smegma buildup underneath the foreskin. This particular penis was left unwashed for 3 days.



Circumcised men can comfortably go several days without washing their genitals, unlike the uncircumcised. Unclean foreskins often lead to infections. Our Asian subject reports that he sometimes goes 2 days without washing his penis. Although Asian foreskins typically produce less smegma than other races, the subject still has experienced infections from his lack of hygiene. Circumcised penises almost never experience infections.